

Pharmasin® 100%

Water Soluble Granules for pigs

- **WATERSOLUBILITY** is crucial to ensure a homogeneous solution
- **ANTIMICROBIAL ACTIVITY** is the real activity against the targeted pathogens
- **STABILITY** is essential for treatment and depending on formulation and packaging



WATERSOLUBILITY

Pharmasin® WSG is formulated to have the best watersolubility and stability


= Correct dose



ANTIMICROBIAL ACTIVITY

Pharmasin® has the highest antimicrobial activity

= Efficacy



STABILITY

Zipper bags for convenient multi-use to guarantee quality

= Stability

Indications

Gastro-intestinal : Ileitis-PIA (*Lawsonia intracellularis*)
Respiratory : *Mycoplasma* spp.

Application

Gastro-intestinal : 5 mg (control) - 10 mg (treatment)
per kg bodyweight for 7 days
Respiratory : 20 mg per kg bodyweight for 10 days



 **HUVEPHARMA®**
We add performance to your business



FOR ANIMAL TREATMENT ONLY
RESTRICTED VETERINARY MEDICINE

Pharmasin[®] 100% Soluble

TYLOSIN TARTRATE WATER SOLUBLE GRANULES FOR USE IN CHICKENS, TURKEYS AND PIGS

Active Ingredient: Tylosin (as tylosin tartrate).....1000 g/kg
Conforming to Ph. Eur.: Tylosin contains not less than 80% Factor A.
Net contents: 1 kg tylosin activity.

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT

INDICATIONS:

Turkeys, broilers & replacement chickens:

As an aid in the prevention and treatment of chronic respiratory disease (CRD) caused by *Mycoplasma gallisepticum* and *Mycoplasma synoviae*. For the treatment of necrotic enteritis caused by *Clostridium perfringens*.

Pigs: Treatment of Porcine Intestinal Adenomatosis (Ileitis) associated with *Lawsonia intracellularis*.

PRECAUTIONS:

Indiscriminate use of this product could contribute to antibiotic resistance. The prophylactic and therapeutic use of this product should only be for the minimum period needed to meet the clinical objective. The product should only be used after consideration of the alternative therapeutic and preventative measures.

MIXING DIRECTIONS:

To assure thorough dissolving, place Pharmasin 100% Soluble (the content of this package) in a mixing container exceeding the pack size, add lukewarm water to the material and stir continuously. Mix this concentrated solution with water and dilute to the required concentration. Always add the water to the powder: do not pour the powder into the water.

Dispose of prepared Pharmasin 100% Soluble solution not used within 24 hours.

DOSAGE:

Poultry: Chronic Respiratory Disease - Mix contents of one container to provide a concentration in drinking water of 0.5 g/L; i.e.

| | | | | |
|-------------------------|-------|-------|---------|----------|
| Pack size | 100 g | 200 g | 1 kg | 25 kg |
| Volume of water treated | 200 L | 400 L | 2,000 L | 50,000 L |

Necrotic Enteritis - Mix contents of one container to provide a concentration in drinking water of 0.05 - 0.15 g/L; i.e.

| | | | | |
|-------------------------|---------------|-----------------|------------------|---------------------|
| Pack size | 100 g | 200 g | 1 kg | 25 kg |
| Volume of water treated | 666 - 2,000 L | 1,333 - 4,000 L | 6,666 - 20,000 L | 166,666 - 500,000 L |

Pigs: Ileitis - Mix contents of one container to provide a concentration in drinking water of 0.1 g/L; i.e.

| | | | | |
|-------------------------|---------|---------|----------|-----------|
| Pack size | 100 g | 200 g | 1 kg | 25 kg |
| Volume of water treated | 1,000 L | 2,000 L | 10,000 L | 250,000 L |

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards.

WITHHOLDING PERIODS:

Meat: Pigs Nil
Poultry Nil
Eggs: Nil

DANGER

Causes serious eye irritation
May cause allergy or asthma symptoms or breathing difficulties if inhaled
May cause an allergic skin reaction
Very toxic to aquatic life

READ LABEL BEFORE USE

Handling precautions:

Wear protective gloves, clothing and eye protection. Wash hands and exposed skin thoroughly after handling
Avoid breathing dust. In case of inadequate ventilation wear respiratory protection.
Contaminated work clothing should not be allowed out of the workplace. Wash contaminated clothing before reuse.

KEEP OUT OF REACH OF CHILDREN

First Aid:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If eye irritation persists: get medical advice/attention.

IF INHALED: If breathing is difficult, remove to fresh air and keep at rest in a position comfortable for breathing.

If experiencing respiratory symptoms: Call a POISON CENTRE or doctor/physician

IF ON SKIN: Wash with plenty of soap and water

If skin irritation or rash occurs: Get medical advice/attention

Environmental Protection:

Avoid release to the environment. Avoid contamination of any water supply with product or empty container. Collect spillage.

Disposal:

Preferably dispose of product by use. Dispose of product and packaging at an approved landfill or equivalent facility.

STORAGE:

Store in original pack, in a dry place, below 25°C.

Protect from light. Do not refrigerate or freeze -protect from frost.

Registered pursuant to the ACVM Act 1997, No. A10039

See www.foodsafety.govt.nz/ for registration conditions.

Registered to: AgriHealth NZ Limited www.agrihealth.co.nz
Manufactured by: Huvepharma AD www.huvepharma.com



HUVEPHARMA[®]

WATER SOLUBLE GRANULES

CHICKENS, TURKEYS AND PIGS

