

# FOR ANIMAL TREATMENT ONLY RESTRICTED VETERINARY MEDICINE Pharmasin® 25% Granular Premix

# TYLOSIN PHOSPHATE FOR USE IN CHICKENS, BEEF CATTLE and PIGS

Active Ingredient: TYLOSIN (as TYLOSIN PHOSPHATE) Tylosin contains not less than 80% factor A Net contents: 20 kg

250 g/kg

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT

INDICATIONS: Beef Cattle:

For the reduction in the incidence of liver abscess.

Broiler Chickens and Replacement Layers:

As an aid in the control of chronic respiratory disease caused by Mycoplasma gallisepticum and/or Mycoplasma synoviae.

For the treatment and prevention of necrotic enteritis associated with susceptible strains of Clostridium perfringens.

Animal Species	Recommended use	Feeding period	Weight of Pharmasin 25% Granular Premix per 1000kg of Complete Feed	Use Level
Pigs	For the treatment and prevention of lleitis Prevention -length of treatment required will be depend on the level of disease Consult your vet for advice	Treatment prevention	400 g 160 g	100 ppm 40 ppm
Broiler Chickens and Replacement Layers	As an aid in the control of chronic respiratory disease caused by Mycoplasma gallisepticum and/or Mycoplasma synoviae.	For 5 days after hatching. Repeat for 2 days at 3 to 5 weeks of age.	3600 g to 4000 g	900 ppm to 1000 ppm
	For the treatment and prevention of necrotic enteritis associated with susceptible strains of <i>Clostridium perfringens</i> .	Treatment -treat for 7 days Prevention -feed during periods of risk of clinical disease	400 g 400 g	100 ppm 100 ppm
Layers	As an aid in the control of chronic respiratory disease caused by Mycoplasma gallisepticum	Feed from time of shifting into laying cages	88 g to 220 g	22 ppm to 55 ppm
Beef Cattle	For reduction in the incidence of liver abscess For supplementary feeds 60-90 mg tylosin/head/day	Feed during period on ration	44 g	11 ppm

The prophylactic and therapeutic use of this product should only be for the minimum period needed to meet the clinical objective. Indiscriminate use of this product could contribute to antibiotic resistance. The product should only be used after consideration of the alternative therapeutic and preventative measures

#### WITHHOLDING PERIODS:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural

Compounds. Meat: Cattle Nil Nil Nil Nil Pigs Poultry Eggs:

## DANGER

Causes serious eye irritation

May cause allergy or asthma symptoms or breathing difficulties if inhaled May cause an allergic skin reaction

Very toxic to aquatic life

# Handling precautions:

Wash hands and exposed skin thoroughly after handling. Contaminated work clothing should not be allowed out of the workplace. Wash contaminated clothing before reuse.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If eye irritation persists: get medical advice/attention.

Avoid breathing dust. In case of inadequate ventilation wear respiratory protection.

IF INHALED: If breathing is difficult, remove to fresh air and keep at rest in a position comfortable for breathing. If experiencing respiratory symptoms: Call the National Poisons Centre 0800 764 766 or a doctor or physician.

IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs: Get medical advice/attention.

## **Environmental Protection:**

Avoid release to the environment of product or empty container. Collect spillage.

Preferably dispose of product by use. Dispose of product and packaging at an approved landfill or equivalent facility.

# UN 3077 ENVIRONMENTALLY HAZARDOUS SUBSTANCE, SOLID, N.O.S. (tylosin phosphate 25%)

#### STORAGE:

Store in original pack, in a dry place, below 25°C. Protect from light. Do not refrigerate or freeze - protect from frost.

Restricted Veterinary Medicine. Registered pursuant to the ACVM Act 1997, No. A010451 See www.foodsafety.govt.nz/ for registration conditions. Approved under the HSNO Act, No. HSR002370. Refer www.epa.govt.nz for controls

Manufactured by: Huvepharma EOOD www.huvepharma.com

Distributed by: AgriHealth NZ Ltd Batch: www.agrihealth.co.nz







Expiry: