



Advanced Non-cycling cow treatment

DIB-Synch Plus



Insemination Timing



All cows showing heat after DIB removal should be mated at the next opportunity.

Note: It is essential for farmers to detect heat and present cows for AI on the two days following DIB removal.

Blanket inseminate all remaining cows, generally within 8 – 20 hours after the final Gonasyn; and 56-72 hours following DIB removal.




Evidence based vet medicines www.agrihealth.co.nz 0800 821 421

AgriHealth


Advanced Heifer/Beef Synchrony



Heifer /Beef Synchrony DIB-CoSynch



Insemination Timing



Set time insemination of all animals in a DIB-CoSynch program typically occurs at the same time as the final Gonosyn injection; and approx. 48 hours following DIB removal.

Further Information

DIB-V is a 1 gram and DIB-h is a 0.5 gram progesterone intravaginal insert registered for use in heifers and cows

The DIB-CoSynch program is designed for simplicity – all injections are a 2mL dose

2mL Gonosyn delivers 100 µg gonadorelin (as acetate)

2mL Cyclase delivers 500 µg cloprostenol (as sodium)

2mL Novormon eCG delivers 400 IU Equine Chorionic Gonadotrophin (eCG)

AI = Artificial Insemination

Gonosyn (A10642), Cyclase (A10490) and Novormon eCG (A10641) are Restricted Veterinary Medicines, available only under veterinary authorisation. DIB-V (A10319) and DIB-h (A10832) are registered Veterinary Medicines. Registered pursuant to the ACVM Act, 1997.



Evidence based vet medicines www.agrihealth.co.nz 0800 821 421

AgriHealth

DIBProgram-ASCARD.7.18