Batch:

Expiry:

Kolibin Neo

Vaccine for cattle containing inactivated Bovine Rotavirus, Bovine Coronavirus, and E. coli.

Contains thiomersal (0.01% w/v)

RESTRICTED VETERINARY MEDICINE
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY



Net Contents 250mL (125 doses)

PAP

109/K/250ml/NZ/1

WARNING

May be harmful if swallowed or inhaled or enters airways. Causes mild skin irritation. May cause an allergic skin reaction.

Handling precautions:

Wear protective gloves when handling.

Wash exposed skin thoroughly after handling.

Avoid breathing mist or spray.

Store locked up.

Contaminated work clothing should not be allowed out of the workplace. Wash contaminated clothing before reuse.

First Aid:

IF SWALLOWED: Immediately call a POISON CENTRE or a doctor/physician. Do not induce vomiting. IF ON SKIN: Wash with plenty of soap and water.

If skin irritation or rash occurs: Get medical advice or attention.

If medical advice is needed, have product container or label at hand.

If exposed or concerned or if you feel unwell: Get medical advice or attention.

For advice call the National Poisons Centre 0800 POISON (0800 764 766).

Disposal:

Any unused product or empty containers should be disposed of at an approved landfill or other approved facility.

Store between 2 - 8°C. Do not freeze. Protect from light. Use within 28 days of opening.

Kolibin Neo is registered pursuant to the ACVM Act 1997, No. A11242.

See www.foodsafety.govt.nz for registration conditions.

Manufactured in Europe by Bioveta, a.s., Ivanovice na Hané, Czech Republic, www.bioveta.cz

bioveta

RESTRICTED VETERINARY MEDICINE
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

Kolibin Neo

Vaccine for cattle containing inactivated Bovine Rotavirus, Bovine Coronavirus, and E. coli.

Contains thiomersal (0.01% w/v)



Net Contents 250mL (125 doses)

8 594004 860583

READ LABEL AND LEAFLET BEFORE USE

Indications

For active immunisation of pregnant heifers and cows to provide calves with passive immunity against gastrointestinal diseases caused by rotavirus, coronavirus and enteropathogenic E. coli strains. When calves consume colostrum and milk from vaccinated cows during the first four weeks of life, these antibodies reduce the incidence and severity of diarrhoea caused by rotavirus, coronavirus and E. coli (K99) and reduce virus shedding by infected calves.

Directions for use

Shake well before use.

CATTLE:

Administer 2mL, by intramuscular injection.
Previously unvaccinated heifers or cows should receive two doses 3 weeks apart with the second dose given 12-2 weeks prior to calving.
Annually revaccinate with a single dose of Kolibin Neo 12-2 weeks prior to each subsequent calving.

Precautions

Do not vaccinate unhealthy animals.

Kolibin Neo can be given as an annual booster vaccination in animals vaccinated with Rotavec Corona (A08312) or ScourGuard 4(K) (A10057) in the previous year.

Withholding Period

Nil

Registered to, and distributed by:
AgriHealth NZ Ltd
0800 821 421
www.agrihealth.co.nz

AgriHealth

Dimensions (mm): **66×66×142**
Colours: **PANTONE**
Pantone 299
Pantone 328
Pantone 369
Pantone Black

Nový Zéland

Pč.:	minimální prostor pro dotisk šarže a exspirace	15×6 mm	ANO / NE	místo pro dotisk je umístěno na největší (nejstabilnější) ploše krabičky	ANO / NE	vyjímku schválil:
Dodavatel:	Dimensions (mm): 66×66×142 Colours: Pantone 299, Pantone 328, Pantone 369, Pantone Black Prepared by: Raška Date: 12. 1. 2016					Approved by: ..... Date: .....