

## Directions:

For adding to the drinking water or liquid diet of calves, pigs and poultry. Must be continuously administered when given in the drinking water, all other sources of water should be removed until the medicated water has been consumed. Prepare fresh solutions daily.

## Dosage:

**Calves:** For bacterial enteritis, salmonellosis, pneumonia, etc. administer approx. 100mg/kg bodyweight in liquid diet or water each day, until 24 hours after symptoms disappear. For a 50kg calf this equates to 5 grams of OxytetSol 10% powder daily.

**Pigs:** For bacterial enteritis, enzootic pneumonia, mastitis, metritis, etc. 2.5 – 4g/litre drinking water daily (approx. 100mg/kg bodyweight daily) until 24 hours after symptoms disappear.

**Poultry:** For respiratory infections, bacterial enteritis, etc. 2.5g/litre drinking water daily (approx. 20mg/kg bodyweight daily) for 5 – 10 days or until 24 hours after symptoms disappear.

Measuring scoop enclosed –  
one level scoop holds 5 grams

AH-OTSP-500g-4

RESTRICTED VETERINARY MEDICINE  
KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY

# OxytetSol

## 10% Powder

Broad spectrum antibiotic soluble powder  
100g/kg Oxytetracycline HCl

**NET WEIGHT 500g**

Please read label completely before use.

OxytetSol 10% Powder is indicated for the treatment and prevention of bacterial infections susceptible to oxytetracycline in calves, pigs and poultry.

## Withholding period:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards. Animals producing meat and offal for human consumption must not be slaughtered during or within 10 days of the last treatment. Not for use in bobby calves. For use in poultry, egg withholding period is 10 days.

Harmful to aquatic life with long lasting effects  
Precautions: Avoid breathing dust. Wear protective gloves after handling.  
First Aid: If medical advice is needed, have product container or label at hand.  
If exposed, or if you feel unwell, call the National Poisons Centre 0800 764 766 or a doctor.  
Environmental protection: Avoid release to the environment of product or empty container.  
Collect spillage of powder or of liquid mixed with powder, and dispose of at an approved facility.

**Disposal:** Shake and empty contents into the medicated water or liquid feed. Preferably dispose of product by use, otherwise dispose of product and empty container at an approved landfill or equivalent facility.

**Storage:** Store in original container in a cool dry place avoiding direct sunlight and temperatures above 30°C.

OxytetSol 10% Powder is registered pursuant to the ACVM Act 1997, No A4144. See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions. **Restricted Veterinary Medicine.**

**AgriHealth**  
Evidence based vet medicines

Registered to AgriHealth NZ Ltd  
[www.agrihealth.co.nz](http://www.agrihealth.co.nz)  
0800 821 421