## FOR ANIMAL TREATMENT ONLY RESTRICTED VETERINARY MEDICINE Pharmasin 100% Soluble

## TYLOSIN TARTRATE WATER SOLUBLE GRANULES FOR USE IN CHICKENS, TURKEYS AND PIGS

Active Ingredient: Tylosin tartrate......1000 g/kg

Conforming to Ph. Eur.: Tylosin contains not less than 80% Factor A.

Net contents: 25 kg tylosin activity.



## READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT INDICATIONS:

Turkeys, broilers & replacement chickens:

As an aid in the prevention and treatment of chronic respiratory disease (CRD) caused by *Mycoplasma gallisepticum* and *Mycoplasma synoviae*. For the treatment of necrotic enteritis caused by *Clostridium perfringens*. Pigs:

Treatment of Porcine Intestinal Adenomatosis (Ileitis) associated with Lawsonia intracellularis.

PRECAUTIONS:

Indiscriminate use of this product could contribute to antibiotic resistance. The prophylactic and therapeutic use of this product should only be for the minimum period needed to meet the clinical objective. The product should only be used after consideration of the alternative therapeutic and preventative measures.

MIXING DIRECTIONS:

To assure thorough dissolving, place Pharmasin 100% Soluble (the content of this package) in a mixing container exceeding the pack size, add lukewarm water to the material and stir continuously. Mix this concentrated solution with water and dilute to the required concentration.

Always add the water to the powder: do not pour the powder into the water.

Dispose of prepared Pharmasin 100% Soluble solution not used within 24 hours.

DOSAGE:

Poultry: Chronic Respiratory Disease - Mix contents of one container to provide a concentration in drinking water of 0.5 q/L; i.e.

Pack size	100 g	200 g	1 kg	25 kg
Volume of water treated	200 L	400 L	2,000 L	50,000 L

Necrotic Enteritis - Mix contents of one container to provide a concentration in drinking water of 0.05-0.15 q/L; i.e.

Pack size	100 g	200 g	1 kg	25 kg
Volume of water treated	666 - 2,000 L	1,333 - 4,000 L	6,666 - 20,000 L	166,666 - 500,000 L

Pigs: Ileitis - Mix contents of one container to provide a concentration in drinking water of 0.1 g/L; i.e.

Pack size	100 g	200 g	1 kg	25 kg
Volume of water treated	1,000 L	2,000 L	10,000 L	250,000 L

It is an offence under the law to sell produce from animals treated with this product with residues that exceed the maximum residue levels in the New Zealand (Maximum Residue of Agricultural Compounds) Food Standards. WITHHOLDING PERIODS:

Meat: Pigs Nil
Poultry Nil

Eggs: Ni

WARNING

Dangerous to the Environment

Handling precautions:

Tylosin may cause respiratory sensitization. Repeated exposure to tylosin may cause skin allergy.

Avoid inhalation and contact with the skin.

KEEP OUT OF REACH OF CHILDREN

**Environmental Protection:** 

Toxic to aquatic organisms. Avoid contamination of any water supply with product or empty container.

Disposal:

Preferably dispose of product by use. Dispose of product and packaging at an approved landfill or equivalent facility.

STORAGE:

Store in original pack, in a dry place, below 25°C. Protect from light. Do not refrigerate or freeze -protect from frost. Registered pursuant to the ACVM Act 1997, No. A10039

See www.nzfsa.govt.nz/acvm/ for registration conditions.

Registered to: AgriHealth NZ Limited <u>www.agrihealth.co.nz</u> Manufactured by: Huvepharma AD <u>www.huvepharma.com</u>

