

**WARNING**  
RESTRICTED VETERINARY MEDICINE  
FOR ANIMAL TREATMENT ONLY  
KEEP OUT OF REACH OF CHILDREN

# Techmulin®

**2.5 % Premix**  
**Tiamulin Hydrogen Fumarate 2.5 %**

Please read label completely before use.

**NET WEIGHT 20 kg**

**Dosage and Administration:**

**Poultry:** Chronic Respiratory Disease:  
**Control:** The dosage of 0.6kg Techmulin 2.5% Premix per 1,000kg of ready-for-use feed administered continuously during laying period.  
**Treatment:** The dosage is 2 kg Techmulin 2.5% Premix per 1,000 kg of ready-for-use feed administered continuously for 5-7 days every 28 days.  
**Mixing directions:** Techmulin 2.5% Premix should be pre-mixed with sufficient carrier to ensure uniform dispersion in the final feed.  
**Warning:** Do not administer to chickens receiving the coccidiostats monensin, narasin or salinomycin.

**Dosage and Administration:**

**Swine:** Swine Enzootic (Mycoplasmal) Pneumonia  
**Control:** (Maintenance of weight gains): During a period of risk, 1.25-1.6 kg of Techmulin 2.5% Premix per 1,000 kg of ready-for use feed administered continuously for 10 days.  
**Prevention:** Swine Dysentery  
The dosage is 1.25-1.6 kg of Techmulin 2.5% Premix per 1,000kg of ready-for-use feed administered continuously during the risk period. Administer to pigs with a history of dysentery but which do not show clinical signs of disease.  
**Treatment:** The dosage is 4kg Techmulin 2.5% Premix per 1,000 kg of ready-for-use feed administered continuously for 5 days.  
**Mixing Directions:** Techmulin 2.5 % Premix should be pre-mixed with sufficient carrier to ensure uniform dispersion in the final feed.  
**Withholding period:** It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (maximum Residue Limits of Agricultural Compounds) Food Standards.  
**Pigs:** producing meat and offal for human consumption must not be slaughtered during or within 5 days of the treatment.  
**Poultry:** producing meat for human consumption must not be slaughtered during or within 3 days of the last treatment.

**Precautions:** May cause organ damage through prolonged or repeated exposure  
Very toxic to aquatic life  
Do not breathe dust. Wear Protective gloves. Wash hands and exposed skin thoroughly after handling.  
**First Aid:** Get medical advice or attention if you feel unwell. If poisoning occurs, call the National Poisons Centre 0800 764 766 or a doctor.  
**Environmental Protection:** Avoid release to the environment of product or empty container. Collect spillage of premix or of feed mixed with premix, use in or as pig or poultry feed.  
**Disposal:** Shake and empty contents into the medicated feed. Preferably dispose of product and empty container at an approved landfill or equivalent facility. Do not use the container for any other purpose.  
**Storage:** Store in original container in a cool dry place avoiding direct sunlight and temperatures above 30°C

Batch No.: Registered to AgriHealth NZ Ltd  
Expiry: 0800 821 421  
www.agrihealth.co.nz

**AgriHealth**



TECHMULIN is registered pursuant to the ACVM Act 1997, No A4209.  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration details. **Restricted Veterinary Medicine.**

Contact the registrant for more information on dosing rates.