FOR ANIMAL TREATMENT ONLY RESTRICTED VETERINARY MEDICINE Pharmasin® 100% Soluble

TYLOSIN TARTRATE WATER SOLUBLE GRANULES FOR USE IN CHICKENS, TURKEYS AND PIGS

Active Ingredient: Tylosin (as tylosin t Conforming to Ph. Eur.: Tylosin contains no Net contents: 1kg tylosin activit READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT

Tylosin (as tylosin tartrate)......1000 g/kg Tylosin contains not less than 80% Factor A. 1 kg tylosin activity.

INDICATIONS:

Turkeys, broilers & replacement chickens:

As an aid in the prevention and treatment of chronic respiratory disease (CRD) caused by Mycoplasma gallisepticum and Mycoplasma synoviae. For the treatment of necrotic enteritis caused by Clostridium perfringens.

Pigs: Treatment of Porcine Intestinal Adenomatosis (Ileitis) associated with Lawsonia intracellularis.

PRECAUTIONS:

Indiscriminate use of this product could contribute to antibiotic resistance. The prophylactic and therapeutic use of this product should only be for the minimum period needed to meet the clinical objective. The product should only be used after consideration of the alternative therapeutic and preventative measures.

MIXING DIRECTIONS:

To assure thorough dissolving, place Pharmasin 100% Soluble (the content of this package) in a mixing container exceeding the pack size, add lukewarm water to the material and stir continuously. Mix this concentrated solution with water and dilute to the required concentration. Always add the water to the powder: do not pour the powder into the water.

Dispose of prepared Pharmasin 100% Soluble solution not used within 24 hours.

DOSAGE:

Poultry: Chronic Respiratory Disease - Mix contents of one container to provide a concentration in drinking water of 0.5 g/L; i.e.

Pack size	100 g	200 g	1 kg	25 kg	
Volume of water treated	200 L	400 L	2,000 L	50,000 L	
Necrotic Enteritis - Mix conter	nts of one contair	ner to provide a o	concentration in	n drinking water	of 0

Necrotic Enteritis - Mix contents of one container to	o provide a concentration in	drinking water of 0.05	- 0.15 g/L; i.e.
---	------------------------------	------------------------	------------------

Pack size	100 g	200 g	1 kg	25 kg
Volume of water treated	666 - 2,000 L	1,333 - 4,000 L	6,666 - 20,000 L	166,666 - 500,000 L
Pigs: Ileitis - Mix contents of one container to provide a concentration in drinking water of 0.1 g/L ; i.e.				
Pack size	100 g	200 g	1 kg	25 kg
Volume of water treated	1,000 L	2,000 L	10,000 L	250,000 L

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: maximum Residue Levels for Agricultural Compounds.

WITHHO	LDING PERIOD	S:
	n '	

meat:	PIQS	NII
	Poultry	Nil
Faas		Nil

DANGER Causes serious eye irritation VATER SOLUBLE GRANULES May cause allergy or asthma symptoms or breathing difficulties if inhaled May cause an allergic skin reaction Verv toxic to aquatic life READ LABEL BEFORE USE Handling precautions: Wear protective gloves, clothing and eye protection. Wash hands and exposed skin thoroughly after handling Avoid breathing dust. In case of inadequate ventilation wear respiratory protection. Contaminated work clothing should not be allowed out of the workplace. Wash contaminated clothing before reuse. KEEP OUT OF REACH OF CHILDREN First Aid: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If eye irritation persists: get medical advice/attention. IF INHALED: If breathing is difficult, remove to fresh air and keep at rest in a position comfortable for breathing. If experiencing respiratory symptoms: Call a POISON CENTRE or doctor/physician IF ON SKIN: Wash with plenty of soap and water If skin irritation or rash occurs: Get medical advice/attention **Environmental Protection:** Avoid release to the environment. Avoid contamination of any water supply with product or empty container. Collect spillage. Disposal Preferably dispose of product by use. Dispose of product and packaging at an approved landfill or equivalent facility. STORAGE Store in original pack, in a dry place, below 25°C. Protect from light. Do not refrigerate or freeze -protect from frost. Registered pursuant to the ACVM Act 1997, No. A010039 See www.foodsafety.govt.nz for registration conditions.

Registered to: AgriHealth NZ Limited www.agrihealth.co.nz Manufactured by: Huyepharma EOOD www.huyepharma.com Agr⁹Health

LB138v5-Phar100wsg1kg-NZ0518