

# Melovem

Higher concentration meloxicam



Evidence based vet medicines

AgriHealth

# Melovem

**HIGHLY POTENT PAIN RELIEF FOR NZ DAIRY COWS**



## Longer-acting NSAID

Single dose provides up to three days of efficacy


## Higher concentration dose

Small volume injection for convenience and improved tolerability

## More doses per bottle

Higher Melovem potency means 50% more cows treated\* per 100mL pack


\* compared with other meloxicam brands



Melovem 100mL - treats 12 cows



Other Meloxicam 100mL - treats 8 cows



# GREATER BENEFITS FOR NZ DAIRY HERDS

Widespread use of Melovem improves health and wellbeing

# Melovem 30

Non-steroidal anti-inflammatory, analgesic and antipyretic injection

For use in cattle to treat pain, inflammation and fever associated with

- acute mastitis
- respiratory infection
- diarrhoea in calves
- disbudding or dehorning

## Directions for Use

**Cattle:** Administer by subcutaneous or intravenous injection. Administer in combination with antibiotic therapy as appropriate. For single use only.

**Dose:** 1.0mL per 60kg body weight (0.5 mg meloxicam/kg).

For young calves a dose rate of 0.3mL per 20kg is appropriate.

## Properties

Meloxicam is a non-steroidal anti-inflammatory drug (NSAID) of the oxicam class which acts by inhibition of prostaglandin synthesis, thereby exerting anti-inflammatory, anti-exudative, analgesic and antipyretic effects.

## Warnings

Do not use in animals suffering from impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of ulcerogenic gastrointestinal disorders, or individual hypersensitivity to the product.

Contraindicated to administer concurrently with glucocorticoids, other non-steroidal anti-inflammatory drugs or with anti-coagulant agents. Should be used with caution in conjunction with other highly protein bound drugs.

Administration is well tolerated in cattle; only a slight transient swelling at the injection site following subcutaneous administration has been observed in some cattle treated in clinical studies.


In the case of overdosage, symptomatic treatment should be initiated.

## Withholding Periods for Cattle

**Meat:** 10 days

**Milk:** 84 hours

Store below 25°C (room temperature). Use remainder of vial within 28 days of opening, or discard unused portion.



Restricted Veterinary Medicine, ACVM Registration Number: A11552. Only available only under veterinary authorisation. Registered to AgriHealth NZ Ltd.



Evidence based vet medicines [www.agrihealth.co.nz](http://www.agrihealth.co.nz) 0800 821 421

**AgriHealth**

MLVM.01.19