

RESTRICTED VETERINARY MEDICINE
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY
CRITICALLY IMPORTANT ANTIBIOTIC

 **TYLOVET® INJECTION**

Tylosin 200mg/mL
solution for injection

250 mL



For intramuscular use in cattle, pigs,
sheep and goats.
For slow intravenous use in cattle.

Read enclosed insert prior to use.

DOSAGE:

Cattle: 5 to 10 mg/kg (2.5 – 5 mL / 100kg) liveweight daily for up to 5 days

Pigs: 5 to 10 mg/kg (0.25 – 0.5 mL / 10kg) liveweight daily for up to 3 days

Sheep and Goats: 10 mg/kg (0.5 mL / 10kg) liveweight daily for up to 5 days

INDICATIONS:

Cattle: Treatment and control of acute mastitis (caused by *Streptococcus uberis*, *S.dysgalactiae*, *S.agalactiae*, *Staphylococcus aureus*, coagulase negative Staphylococci), metritis caused by *Trueperella pyogenes*, respiratory infections caused by *Pasteurella multocida*, foot-rot (necrotic pododermatitis) and calf diphtheria caused by *Fusobacterium necrophorum*.

Pigs: Treatment and control of ileitis associated with *Lawsonia intracellularis*, swine erysipelas caused by *Erysipelothrix rhusiopathiae*, pneumonia caused by *Mycoplasma hyopneumoniae* and *M. hyorhinis*, and arthritis due to *Mycoplasma hyosynoviae* and *M. hyorhinis*.

Sheep and Goats: Treatment of the early stages of peracute and acute contagious agalactia caused by *Mycoplasma agalactiae* and caprine pleuropneumonia caused by *Mycoplasma mycoides* var *capri* (*M.capri*).

The prophylactic and therapeutic use of this product should only be for the minimum period needed to meet the clinical objective. Indiscriminate use of this product could contribute to antibiotic resistance.

 **AgriHealth**

Produced in Europe by Huvepharma E00D
Registered to and distributed by
AgriHealth NZ Ltd
www.agrihealth.co.nz, Phone 0800 821 421
Registered pursuant to the ACVM Act 1997, No. A10807. See foodsafety.govt.nz for registration conditions



SOLUTION FOR INJECTION

CATTLE, SHEEP, GOATS, SWINE

RESTRICTED VETERINARY MEDICINE
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY
CRITICALLY IMPORTANT ANTIBIOTIC

 **TYLOVET® INJECTION**

Tylosin 200mg/mL
solution for injection

250 mL



For intramuscular use in cattle, pigs,
sheep and goats.
For slow intravenous use in cattle.

WARNING

May cause an allergic skin reaction

Causes eye irritation

Safety precautions and directions

Wear protective gloves. Wash hands thoroughly after handling. Contaminated clothing should not be allowed out of the workplace. Wash contaminated clothing before reuse.

First Aid

If in the eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists, get medical advice.

If on skin: Wash with plenty of soap and water. If skin irritation or rash occurs, get medical advice. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor.

WITHHOLDING PERIODS:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Meat: Animals producing meat and offal for human consumption must not be sold for slaughter either during treatment or within 21 days of the last treatment.

Milk: Milk intended for sale for human consumption or manufacture for human consumption must be discarded during treatment and for 72 hours following the last treatment.

STORE below 25° C, protect from light.
Discard unused content 28 days after opening.



SOLUTION FOR INJECTION

CATTLE, SHEEP, GOATS, SWINE