



Hostazym[®] X

LB1559v6HostX50fa-20-NZ1124

MICROGRANULATE

Inclusion Rate 50 g/ton
ENZYME FEED ADDITIVE

PRODUCT DESCRIPTION

An enzyme preparation of endo-1,4-beta-xylanase (EC 3.2.1.8) produced by *Trichoderma sp.* having a minimum activity of 30,000 EPU/g.

RECOMMENDED INCLUSION RATE

For use in monogastric animal feed rich in non-starch polysaccharides.

The recommended doses are as follows:

Species or category of animal	Units of activity per kg of complete feed		Recommended inclusion rate of Hostazym [®] X, g/ton of feed
	Minimum activity per kg of complete feed	Maximal activity per kg of complete feed	
Chickens for fattening	1500 EPU	-	50
Turkeys for fattening	1050 EPU	-	35

Laying hens	1500 EPU	-	50
Piglets	1500 EPU	-	50
Pigs for fattening	1500 EPU	-	50

DIRECTIONS FOR USE

To guarantee stability for a minimum of 24 months, Hostazym[®] X must be stored unopened in the original packaging in a dry place at room temperature.

Temperatures above 85°C during pelleting may reduce the enzymatic activity.

DANGER

May cause allergy or asthma symptoms or breathing difficulty if inhaled.

HANDLING PRECAUTIONS:

Avoid breathing dust. In case of inadequate ventilation, wear respiratory protection.

FIRST AID: IF INHALED: remove person to fresh air and keep comfortable for breathing. If experiencing respiratory symptoms, call the National Poisons Centre 0800 764 766 or a doctor.

DISPOSAL: Preferably dispose of product by use. Shake empty bag into medicated feed. Dispose of unused product and empty packaging at an approved landfill or other facility.

Batch:

Mfg. date:

Exp. date:

Net weight 20 kg

Gross weight 20.5 kg



3 800043 740195



C/PAP/PE



MANUFACTURER

HUVEPHARMA EOOD
www.huvepharma.com

DISTRIBUTOR

AgriHealth NZ Limited
www.agrihealth.co.nz
Phone: 0800 821 421

AgriHealth