



PENETHAMATE HYDRIODIDE
PenethaOne[®]
Antibiotic Injection

**RESTRICTED VETERINARY MEDICINE
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY**

PenethaOne[®]
Antibiotic Injection
PENETHAMATE HYDRIODIDE

For the treatment of mastitis, uterine and respiratory infections and footrot in cattle, caused by penicillin-sensitive Gram-positive bacteria.

Highly important antibiotic

1 x 10 MIU vial antibiotic
1 x 36 mL vial diluent

**READ THE ENCLOSED LEAFLET
BEFORE USING THIS PRODUCT**

INDICATIONS:

PenethaOne is indicated for the treatment and control of infections caused by penicillin-sensitive gram-positive bacteria that cause mastitis, uterine and respiratory infections and footrot in cattle.

DIRECTIONS FOR USE:

Reconstitute the powder contained in the vial by adding all the contents (36mL) of the diluent supplied. Shake well after reconstitution.

Administer by intramuscular (IM) injection only, in the anterior half of the neck. The reconstituted aqueous suspension is stable for 1 day at 25°C (room temperature) or for 7 days when stored between 2 - 8°C (refrigerated). Use aseptic technique when withdrawing and administering each dose of PenethaOne.

Dosage: 1. Cattle: 5 MIU once daily (equivalent to 10,000 - 15,000 IU/kg bodyweight/day). Repeat dose daily for 1 - 5 days. **2.** For mastitis in lactating dairy cattle: Use the above dosing or alternatively give 10 MIU on day one, followed by 5 MIU on day two.

A maximum dose volume of 20mL per injection site should be administered.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIOD: Not for use in bobby calves. **MEAT (Cattle):** Animals producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 7 days of the last treatment.

MILK (Cattle): Milk intended for sale for human consumption must be discarded during treatment and for not less than: 48 hours after the last treatment of 5 MIU daily dosage; or 60 hours after the last treatment of 10 MIU plus 5 MIU.

Contraindications:
DO NOT administer by intravenous (IV) injection.
DO NOT use in animals that are known to be hypersensitive to penicillins or cephalosporins.
DO NOT administer to animals with renal disease including anuria or oliguria.
DO NOT administer to lagomorphs or small rodents.

Adverse Reactions:
In very rare cases, anaphylactic shock may occur.

STORAGE:
Store below 25°C. Keep away from direct sunlight.

Registered pursuant to the ACVM Act 1997, No A11828
See www.foodsafety.govt.nz for registration conditions.

Registered to:
Divasa-Farmavic, S.A.
Spain



AgriHealth



Distributed by:
AgriHealth NZ Ltd
www.agrihealth.co.nz
Phone 0800 821 421



BN:
EXP:

9110851000-0