MeloxiVet



AgriHealth

RESTRICTED VETERINARY MEDICINE KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY Meloxicam 20mg/mL

Non-steroidal anti-inflammatory, analgesic and antipyretic injection for use in cattle, pigs and horses.

Withholding Period:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residues Limits of Agricultural Compounds) Food Standards.

Meat: Cattle-10 days Pigs-3 days Horses-28 days Milk: Cattle-84 hours

Store below 25°C. Use within 28 days of opening.

READ CARTON LABEL AND LEAFLET BEFORE USE

Directions for use: Administer in combination with antibiotic therapy as appropriate.

CATTLE: SINGLE USE ONLY. Administer 2.5mL per 100kg body weight (0.5 mg meloxicam /kg) by subcutaneous or intravenous injection. For very young calves weighing less than 50kg a dose rate of 0.5mL per 20kg is appropriate.

PIGS: Administer 2mL per 100kg bodyweight by intramuscular injection in the anterior neck (0.4 mg meloxicam /kg). Can be repeated once after 24 hours if necessary.

HORSES: SINGLE USE ONLY. Administer 3mL per 100kg body weight by intravenous injection (0.6 mg meloxicam /kg).

I Δ10676

Ratch:

Expiry:

Manufactured in Europe. Registered in New Zealand to: AgriHealth NZ Ltd 0800 821 421 www.agrihealth.co.nz

