

FOR ANIMAL TREATMENT ONLY
KEEP OUT OF REACH OF CHILDREN

NicarbMax 100%

For the prevention and control of coccidiosis caused by
Eimeria acervulina, *E. maxima*, *E. mivati*, *E. necatrix*, and *E. tenella*
in broiler chickens

Active Ingredient: Nicarbazine 1,000 g / kg

Net Contents : 20kg

Mixing Directions:

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Mix 125g of NicarbMax 100% per tonne of feed to produce a concentration of 125 g / tonne nicarbazine in the finished broiler feed.

NicarbMax 100% must be thoroughly mixed in the feed to assure even distribution. It is recommended that the required quantity of NicarbMax 100% is pre-blended with 4-5kg of ground cereal before adding to the total mix.

Feed containing NicarbMax 100% should be the only ration fed to broiler chickens during treatment. Use cautiously for 14 days, or as directed by a veterinarian. **Do not use in layer hens.** Consult a veterinarian for advice relating to coccidia control.

Contraindication:

DO NOT USE in birds which are producing or may in the future produce eggs or egg products for human consumption as egg quality and production may be affected.

Precaution:

Use in birds greater than 28 days of age may increase susceptibility to heat stress.

WITHHOLDING PERIODS:

Meat: NIL

Eggs: DO NOT USE in birds that are producing or may in the future produce eggs or egg products which may be used or processed for human consumption.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Limits for Agricultural Compounds.

| | |
|------------|--|
| STORAGE: | Store in a dry place below 25°C. Protect from light and moisture. |
| DISPOSAL: | Shake and empty containers into medicated feed. Do not dispose of unused product on-site. Puncture or shred and bury empty container in a site specifically designed for the purpose away from waterways, vegetation and roots. Empty bag and left-over product should not be burnt. |
| FIRST AID: | IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs seek medical advice. Wash contaminated clothing before reuse. |

Registered to: AgriHealth NZ Ltd
www.agrihealth.co.nz

AgriHealth

Registered pursuant to the ACVM Act, 1997, No 10785.
See www.foodsafety.govt.nz for registration conditions

BATCH:

EXPIRY:



9 421906 196092 >

AH-NM-20kg-2