

Keep out of the reach of children
FOR ANIMAL TREATMENT ONLY

DIB-h

Intravaginal Insert containing 0.5 gram
progesterone per device

NET CONTENTS

10 Intravaginal Inserts

Each insert contains 0.5 gram progesterone



DIB-h is an intravaginal progesterone release insert for controlled breeding in cattle.

Each DIB-h insert contains 0.5 gram progesterone in an inert silicone elastomer.

INDICATIONS

- Treatment of post-partum anoestrous (non-cycling) cows.
- Control of the oestrous cycle in maiden heifers and cows.
- To enable synchronisation for:
 - concentrated artificial insemination (AI).
 - embryo collection and transfer programs.

DIRECTIONS FOR USE

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

DOSE

One DIB-h insert per animal.

TREATMENT PROGRAMS

DIB-h has been shown to be effective for cattle reproduction synchrony programs when inserted for a period of between 5 and 8 days. For best results, DIB-h is recommended for use as part of a structured synchrony program in conjunction with Gonasyn (A10642) and Cyclase (A10490) in synchrony programs, and also Novormon (A10641) in synchrony programs for non-cycling cows. Consult your veterinarian for further information, as reproductive treatment and oestrus synchrony programs may be modified according to veterinary advice.

Insertion

1. Wear rubber gloves when handling inserts.
2. Wash the applicator in a non-irritating antiseptic solution and dip the front portion in veterinary obstetrical lubricant to assist ease of insertion.
3. Place the insert into the applicator so that the wings are folded and the tips are protruding from the front of the applicator.
4. Dip the front of the applicator into veterinary obstetrical lubricant.
5. Lift the cow's tail and wipe the vulva clean.
6. With tail lifted insert the loaded device, sloping it slightly upwards, through the vulva and then forward, into the forward portion of the vagina. NB Do not insert into overly small animals when application discomfort is evident.
7. Release the insert from the applicator and then remove the applicator.
8. Approx. 10 cm of the tail of the insert should be visibly protruding from the vulva.
9. Ideally mark each treated animal to easily identify for removal of the insert and DIB-Synch program completion.

Removal

Remove by gently, but firmly, pulling on the protruding tail of the insert. If no tail is obvious, conduct a vaginal examination of the cow to check for the insert (then remove).

Contraindications

Not suitable for use in animals with abnormal vaginas. Animals in poor condition resulting from illness or nutritional stress may not respond.

Withholding Period

Meat: nil

Milk: nil

Remove inserts before slaughter.

WARNING

Suspected of damaging fertility or the unborn child.

Suspected of causing cancer.

Obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

Wear protective gloves.

If exposed or concerned: Get medical advice/attention. For advice contact the National Poisons Centre 0800 POISON (0800 764 766).



Disposal

Used inserts and packaging should be burnt in an appropriate incinerator, buried in an approved landfill, or disposed of by approved methods.

Store locked up at room temperature, out of reach of children.

DIB-h is registered pursuant to the ACVM Act 1997, N° A10832. See www.foodsafety.govt.nz for registration conditions.

MANUFACTURER:

SYNTEX S.A.

www.syntexar.com

REGISTERED TO:

AgriHealth

AgriHealth NZ Ltd
www.agrihealth.co.nz
Phone 0800 821 421

Batch N°/Manufacture Date/Expiry Date: See bag.



09421906196023