



C1477-617NZ/05-14

EX2888-16-02

Warnings

Do not use concurrently with another non-steroidal anti-inflammatory. Do not administer where there is abundant haemorrhage. Do not use in animals suffering from cardiac, hepatic or renal disease. Do not administer by the intra-arterial route. Do not administer to an animal that has developed hypersensitivity to ketoprofen. Abomasal ulceration and perforation has been seen when used in calves at disbudding.

Handling Precautions

KEEP OUT OF REACH OF CHILDREN

WARNING: Harmful if swallowed. Causes skin and eye irritation. Harmful to terrestrial vertebrates. Wash hands and exposed skin thoroughly after handling. Do not eat, drink or smoke when using this product. Wear protective gloves. Take off contaminated clothing and wash before reuse. If medical advice is needed, have product container or label at hand.

IF SWALLOWED: Call the National Poisons Centre 0800 764 766 or a doctor/physician if you feel unwell. Rinse mouth.

IF ON SKIN: Wash with plenty of soap and water. If skin irritation occurs: Get medical advice / attention.


IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.

DISPOSE of empty container by wrapping with paper and putting in the garbage. Avoid release of the product into the environment.

KetoMax 15% is registered pursuant to the ACVM Act 1997, No. A11031. See www.foodsafety.govt.nz for registration conditions.

**FOR ANIMAL TREATMENT ONLY
RESTRICTED VETERINARY MEDICINE**

KetoMax 15%



For use in cattle for the anti-inflammatory, analgesic and anti-pyretic treatment of:

- Acute mastitis
- Respiratory conditions
- Musculoskeletal conditions
- Mammary oedema
- Colic

READ THE LABEL BEFORE USING THIS PRODUCT

Directions for Use

CATTLE: Administer by intravenous or intramuscular injection, in the neck.
DOSE: 1 mL / 50 kg / day (3 mg/kg ketoprofen) for 1 to 3 days depending on severity and duration of symptoms

Properties

Ketoprofen is a non-steroidal anti-inflammatory drug (NSAID) of the propionic acid class. The anti-inflammatory, analgesic and anti-pyretic effects of ketoprofen are attributed to the inhibition of cyclooxygenase in the inflammatory process. Ketoprofen is rapidly and extensively absorbed via either intramuscular or intravenous injection.

Withholding Periods

Meat: Cattle producing meat or offal for human consumption must not be sold for slaughter either during treatment or within **2 days** of the last treatment.


Milk: Nil

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residues Limits of Agricultural Compounds) Food Standards. Store below 25°C (room temperature). Use remainder of vial within 28 days of opening, or discard unused portion.

**Registered to AgriHealth NZ Ltd
Freephone 0800 821 421
www.agrihealth.co.nz**

**FOR ANIMAL TREATMENT ONLY
RESTRICTED VETERINARY MEDICINE**

KetoMax 15%



Batch

Expiry



Descripción / Description	CTN KetoMax 15% 250ml NZ	Date 16/08/2016	Proof 2
Número Producto / Item Number	EX2888-16-02	Tintas / Inks	Colores Técnicos / Technical Colours
Plano / Keyline		P369	P471 Black Keyline
Tamaño / Size			
Gramaje / Weight			
Códigos / Codes			
Texto Regulatorio / Regulatory text			
Technical Approval	Regulatory Affairs	Marketing	Manufacturing
Date: Observations:	Date: Observations:	Date: Observations:	Date: Observations:
Fuentes / Fonts:	Artworker: ThinkInpress AMS Project: 16ECU0103		