

**KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY**

STENOROL®

Premix for Poultry

Net contents: 20 kg

Active ingredient: HALOFUGINONE HYDROBROMIDE 6 g/kg

READ THE ENTIRE LABEL BEFORE USE

INDICATIONS:

As an aid in the prevention of coccidiosis caused by *Eimeria acervulina*, *E. maxima*, *E. necatrix*, and *E. tenella* in broiler chickens.

DIRECTIONS FOR USE:

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

POULTRY: For use in broiler chickens. STENOROL should be mixed into a secondary premix prior to incorporation in finished feed. Add 0.5kg of STENOROL to each tonne of chicken feed. This produces a halofuginone hydrobromide content of 3g/tonne in the finished feed which should be fed continuously as the sole ration.

NOTE: Medicated feed must be labelled with appropriate precautions.

CONTRAINDICATIONS:

The use of this additive is not recommended for young quail, pigeons, red-legged partridges, young guinea fowl, ducks or water fowl.

WITHHOLDING PERIOD:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Meat: Animals producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 2 days of the last treatment.

STORAGE:

Store below 25°C in a dry place.

WARNING

Harmful if swallowed. May cause an allergic skin reaction.

Toxic to aquatic life with long lasting effects. Toxic to terrestrial vertebrates.

Handling precautions:

This product must only be used as a veterinary medicine applied to poultry feed. Wear protective gloves, face protection and coveralls buttoned to the neck when handling. Do not eat drink or smoke when using this product. Wash hands and exposed skin thoroughly after handling. Take off contaminated clothing and wash before reuse. Contaminated work clothing should not be allowed out of the workplace. Do not breathe dust.

First Aid:

IF medical advice is needed, have product container or label at hand.

IF SWALLOWED: Rinse mouth. Call the National Poisons Centre (0800 POISON, 0800 764 766) or a doctor if you feel unwell.

IF ON SKIN: Wash with plenty of soap and water. If skin rash or irritation occurs, get medical advice or attention.

Environmental protection:

Avoid release to the environment. Collect spillage.

The active ingredient in this substance is persistent in the environment. The maximum application rate of manure from poultry treated with Stenorol that should be applied to land is equivalent to 170 kg N/ha per annum to ensure that the active ingredient does not build up in soil. Stenorol or manure from poultry treated with Stenorol should not be directly applied into or onto water.

Disposal:

Preferably dispose of product by use. Shake bag empty into medicated feed. Dispose of unused product and empty packaging at an approved landfill or other facility.

BATCH:

Manufactured in Europe by:
HUVEPHARMA EOOD
www.huvepharma.com

Registered pursuant to the ACVM Act 1997, No A11396.
See www.foodsafety.govt.nz for registration conditions.
Registered to: AgriHealth NZ Ltd
www.agrihealth.co.nz

EXPIRY: