# FACT SHEET Toltrox



## WHAT IS TOLTROX?

An oral suspension for the treatment and prevention of coccidiosis in calves, containing Toltrazuril 50mg/mL

## WHAT IS TOLTROX FOR?

Toltrox controls coccidiosis caused by *Eimeria bovis* or *Eimeria zuernii* in young cattle (calves).

## WHEN SHOULD YOU USE TOLTROX?

Coccidiosis typically occurs in cattle less than a year old causing diarrhoea which is often blood stained. Calves recently weaned from meal are particularly susceptible to outbreaks of clinical disease. Toltrox is effective in one dose because it attacks all stages of the coccidia parasite in the calf.

Farmers should obtain veterinary advice regarding the cause of calf scours, and subsequent actions to mitigate future deleterious impacts. Treatment of calves at the time of meal withdrawal will control coccidiosis associated with weaning.

Treatment of all calves on the farm as soon as clinical signs of disease are seen in herd mates will reduce the impact of a coccidiosis outbreak. A single treatment rapidly acts to stop further intestinal damage in affected animals, and prevents herd mates from developing diarrhoea.

To obtain maximum benefit on farms with a history of coccidiosis, Toltrox should be given approximately one week prior to the expected onset of clinical signs, or at weaning from meal.

## **HOW DO YOU USE TOLTROX?**

Administer 3mL Toltrox per 10kg bodyweight orally (i.e. 30mL per 100kg calf)

## WITHHOLDING PERIOD

Cattle: Meat 56 days

## **STORAGE**

Store below 25°C in original container

## **PACK SIZES**

Available in 1 litre and 5 litre plastic backpack bottles

REGISTERED VETERINARY MEDICINE ACVM Registration Number: A11401



**FS-TLX 2020** 

AgriHealth