

Active Constituents: This vaccine contains viable oocysts of *Eimeria acervulina* Strain RA₃₊₂₀, *E. maxima* Strain MCK₊₁₀, *E. necatrix* Strain mednec₃₊₈ and *E. tenella* Strain Rt₃₊₁₅ suspended in phosphate buffered saline (PBS). Each dose comprises a minimum of *E. acervulina* 50 oocysts, *E. maxima* 100 oocysts, *E. necatrix* 100 oocysts and *E. tenella* 150 oocysts, with a minimum predicted titre of 1.6×10^4 oocysts per mL at the end of the shelf-life.

Statement of Claims: Live oocyst preparation to aid in the control of the four major species of *Eimeria* that cause Coccidiosis in breeders, broilers and layer chickens.

1000 Doses

Contents: 1000 doses in a plastic bottle. The *Eimeria* oocysts are supplied as a suspension of oocysts in 25 mL of PBS contained within a single capped bottle.

READ DIRECTIONS CAREFULLY

DIRECTIONS FOR USE

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Contraindications:

- This product is contraindicated for vaccinating unhealthy birds.

Precautions:

- Do not use in-feed or water anticoccidial medications at or soon after the time of vaccination.
- Avoid administration of sulfa drugs immediately following vaccination with EIMERIAVAX 4m as this will delay oocyst output and impair immune response.
- Avoid vaccinating with EIMERIAVAX 4m unless birds are protected from exposure to litter from the previous batch for up to 21 days post-vaccination.
- Do not use EIMERIAVAX 4m as a diluent for other medications or treatments.
- Avoid using in birds in lay.

Dosage and Administration

Eyedrop Method of Administration:

USE ALL PRODUCT WITHIN 6 HOURS OF OPENING.

Birds from day-old can be vaccinated with EIMERIAVAX 4m. With individual bird administration, it is essential for each bird to receive a full dose. A second dose should be administered if the first is not fully absorbed by the eye. It is recommended that all birds be vaccinated prior to 18 weeks of age. The most appropriate age to vaccinate may be determined from knowledge of the challenge conditions applicable to the flocks concerned. The effect of EIMERIAVAX 4m vaccination during lay has not been assessed.

Preparation of vaccine for use and general instructions:

- Prior to use gently shake vials to fully resuspend all oocysts.
- Remove the cap and insert an eye-dropper tip or other administration device, calibrated so as to distribute 25 microlitre drops of vaccine.
- Hold each bird with its head tilted to one side.
- If using a dropper tip attached to the bottle, invert the dropper bottle.
- Allow a single drop to form at the tip and fall freely into the open eye, gently flooding it. The drop (before release) and dropper tip or administration device should NOT touch the eye surface.
- Allow the bird to blink before releasing it.
- Administer a second drop if the bird did not receive a full dose.

Spray method of administration

EIMERIAVAX 4m has been demonstrated effective when administered using a hatchery spray apparatus that delivers 24 mL of diluted vaccine per box of 100 day-old chicks, i.e. 0.24 mL per chick. Please note the following points:

- The vaccine is to be diluted in water. Blue food grade dye should be added to the diluent prior to spraying to encourage uptake of the vaccine by preening. If using a spray apparatus that delivers 24 mL per 100-chick box, one bottle of EIMERIAVAX 4m is diluted to a total volume of 240 mL.

- Open the vaccine bottle when required by peeling back the foil cap and removing the rubber bung. Rinse the vaccine bottle several times in the water diluent to ensure all the oocysts are removed from the bottle.
- Ensure the diluted vaccine is regularly agitated during the spray applications to ensure vaccine oocysts do not settle.
- The spray unit should deliver an even mist with the added blue dye evident. Ensure mist covers all chicks.
- After spraying, leave the chicks in a light place to allow preening and vaccine ingestion.
- Use all vaccine within 6 hours of dilution.
- Please contact **AgriHealth NZ** Ltd if further information regarding specific spray units is required.

Bird management after administration

- Continued exposure to oocysts shed in faeces is essential for the development of an effective immune response.
- Birds can be vaccinated from day-old.
- Anticoccidial medication is never recommended.
- If depression associated with vaccination is noted, light intensity should be increased and/or extended and extra feed provided.
- Monitor birds carefully for 4 weeks post-vaccination. Some lesions can be expected. Lesions can be associated with an immune response and should not be used as the sole criterion for assessing efficacy.
- Investigate all suspect coccidiosis lesions as this vaccine does not protect against *Eimeria* species that are not represented in the vaccine. Monitor gross lesion scores for 4 weeks post-vaccination.

Duration of immunity: Continuing exposure to oocysts in the environment is essential and provides ongoing stimulus to the immune system. Under these exposure conditions, a protective immune response beyond 18 weeks of age may be experienced. When birds are removed from the source of vaccine oocysts, such as when they are transferred between facilities, immunity may wane due to reduced exposure to cycling oocysts. As a result, birds may subsequently become susceptible to field infections from *Eimeria* species represented in the vaccine.

WITHHOLDING PERIOD: NIL

USER SAFETY INFORMATION:

Accidental contact of EIMERIAVAX 4m with eyes or mouth may cause irritation. Eyes should be rinsed immediately and medical advice sought. The mouth should be rinsed two or three times with a mouth gargle. Additional information is available on the product Safety Data Sheet.

FIRST AID:

If poisoning occurs, contact a Doctor.

DISPOSAL:

Discard unused vaccine and empty vaccine bottles into a disinfectant solution (e.g. chlorine-based bleach). Dispose of any such inactivated, unused vaccine and empty containers by wrapping in paper and putting in garbage. In cases of spillage, soak up the liquid with an absorbant sponge or cloth and incinerate. Treat the surface with a disinfectant solution (e.g. chlorine-based bleach).

CAUTION:

The capability of this vaccine to produce satisfactory results depends upon many factors including, but not limited to, conditions of storage and handling by the user, administration of the vaccine, health and responsiveness of individual birds and degree of field exposure. Therefore, directions for use should be followed carefully.

STORAGE INSTRUCTIONS:

Product should be stored between 2 °C and 8 °C (refrigerated) away from sunlight. Do not freeze. Product should be used by the expiry date. Use all product within 6 hours of opening and dilution.

CONTACT DETAILS:

Registered to and distributed by:

AgriHealth NZ Ltd
Ph 0800 821 421
www.agrihealth.co.nz



Manufactured in Australia by:

BIOPROPERTIES Pty Ltd
for EIMERIA Pty Ltd



AgriHealth NZ Ltd markets and sells EIMERIAVAX 4m on behalf of Huvepharma EOOD.

Eimeriavax 4m is registered pursuant to the ACVM Act 1997, No. A11638 See www.foodsafety.govt.nz for registration conditions.