



Injectable solution

Cyclase®
100mL

Manufactured by
Syntex S.A.
www.syntexar.com

Distributed by

AgriHealth

Agrihealth NZ Ltd
www.agrihealth.co.nz

Restricted Veterinary Medicine

Registered pursuant to the ACVM Act 1997
Nº A10490
See www.foodsafety.govt.nz for registration conditions

Approved under the HSNO Act,
Nº HSR002169
See www.epa.govt.nz for controls

RESTRICTED VETERINARY MEDICINE
KEEP OUT OF THE REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

Cyclase®
100mL

Injectable solution

Contains:
Cloprostenol (as sodium)
250 µg/mL

For luteolysis of functional corpora lutea
in cows, pigs and horses

Complements use of the DIB-V (A10319)
and DIB-h (A10832) progesterone inserts
and Gonasyn (A10642) in DIB-Synch and
DIB-CoSynch oestrus synchrony
programs, and also Novormon eCG
(A10641) in DIB-Synch Plus programs.
Can be used as a diluent
for Novormon eCG.



INDICATIONS

For luteolysis of functional corpora lutea in cows, sows and mares. Manipulation of oestrus cycles, treatment of retained foetal membranes, pyometra or chronic endometritis and induction of abortion and parturition in cows.

DOSEAGE:

In DIB-Synch and DIB-Synch Plus programs in cattle: 2mL

ADMINISTRATION

Administer by intramuscular or subcutaneous injection in the anterior half of the neck. Read the enclosed leaflet before use.

CONTRAINDICATIONS

Do not use in pregnant animals unless the objective is to terminate pregnancy. Do not administer IV.

PRECAUTIONS:

Cloprostenol may damage fertility or the unborn child. May cause respiratory problems including bronchospasm in high doses. Cloprostenol can be absorbed through the skin, avoid skin contact and exercise extreme caution especially in women of child bearing age and people with bronchial disease such as asthmatics. Obtain special instructions before use, and do not handle until all safety precautions have been read and understood. Use protective personal equipment as required to reduce risk of skin contact and inhalation.

Do not breathe mist or spray. Do not eat, drink or smoke when using this product. Wash hands and exposed skin thoroughly after handling.

FIRST AID

Wash exposed skin immediately with plenty of water and soap. If inhaled move to fresh air. If exposed or concerned, seek medical advice or attention. For Advice contact the National Poisons Centre 0800 764 766 or seek Medical Advice from a Doctor.

DISPOSAL

Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility.

STORAGE

Store below 25°C away from direct light. Store locked up. Use within 30 days of opening the pack. When used as a diluent for Novormon eCG (A10641), store refrigerated (2-8°C) and discard contents 21 days after mixing.

RESTRICTED VETERINARY MEDICINE
KEEP OUT OF THE REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

Cyclase®
100mL

Injectable solution

Contains:
Cloprostenol (as sodium)
250 µg/mL

For luteolysis of functional corpora lutea
in cows, pigs and horses

Complements use of the DIB-V (A10319)
and DIB-h (A10832) progesterone inserts
and Gonasyn (A10642) in DIB-Synch and
DIB-CoSynch oestrus synchrony
programs, and also Novormon eCG
(A10641) in DIB-Synch Plus programs.
Can be used as a diluent
for Novormon eCG.



CELLIP 19/2

BATCH N°:

EXPIRY DATE: