

RESTRICTED VETERINARY MEDICINE
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

BaciMax 150 Granular

Zinc Bacitracin

For the prevention and treatment of necrotic enteritis
(*Clostridium perfringens* types A and C) in pigs and poultry

Active Ingredient: Zinc Bacitracin to provide 150 g per kg bacitracin
(1g bacitracin is equivalent to 42,000IU)

Net Contents : 25kg

Mixing Directions: BaciMax 150 Granular should be premixed with 10 - 20kg of feed before being thoroughly mixed with bulk feed.


Species	kg BaciMax 150 Granular / tonne feed	equivalent to bacitracin ppm	Indications for Use
Broiler Chickens	0.33	50	As an aid in the prevention of necrotic enteritis caused by or complicated by <i>Clostridium</i> spp. or other organism susceptible to zinc bacitracin.
	1.33	200	As an aid in the treatment of necrotic enteritis caused by or complicated by <i>Clostridium</i> spp. or other organism susceptible to zinc bacitracin. CAUTION: To control a necrotic enteritis outbreak, start medication at the first clinical signs of disease. Administer continuously for 5 - 7 days or as long as clinical signs persist and then reduce medication to preventative level (0.33 kg BaciMax 150 Granular / tonne feed).
Quail	1.33	200	For the prevention of ulcerative enteritis in growing quail due to <i>Clostridium colinum</i> susceptible to zinc bacitracin. Feed continuously as the sole ration.
Pregnant sows	1.65	250	For the control of clostridial enteritis caused by <i>Clostridium perfringens</i> in suckling piglets when fed to sows from 14 days before through to 21 days after farrowing on premises with a history of clostridial scours. CAUTION: Diagnosis should be confirmed by a veterinarian when results are not satisfactory. Feed containing an approved level of zinc bacitracin should be the sole ration.

Caution:

Indiscriminate use may increase the risk of the development of antibiotic resistance. Duration of use should be limited to risk periods when possible.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIOD: NIL

WARNING:	May cause an allergic skin reaction.	
PRECAUTIONS:	Read label before use. Avoid breathing dust. Contaminated work clothing should not be allowed out of the workplace. Wash contaminated clothing before reuse. Wear protective gloves, clothing and face protection.	
STORAGE:	Store in a dry place below 25°C. Close container securely after use.	
DISPOSAL:	Shake and empty containers into medicated feed. Do not dispose of undiluted chemicals on-site. Puncture or shred and bury empty container in a site specifically designed for the purpose away from waterways, vegetation and roots. Empty bag and left-over product should not be burnt.	
FIRST AID:	IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs seek medical advice or attention.	

Batch:	
Expiry:	

Registered to: AgriHealth NZ Ltd
www.agrihealth.co.nz

Registered pursuant to the ACVM Act, 1997 No A10742.
See www.foodsafety.govt.nz for registration conditions





AH-BM-25kg-3