

AgriHealth
Evidence based vet medicines

An aid in the prevention of coccidiosis in broiler chickens caused by *Eimeria maxima*, *E. tenella* and *E. acervulina*.

10 X 10,000 Dose Vials

Coccidiosis vaccine containing live oocysts of *Eimeria maxima*, *E. tenella* and *E. acervulina*

ADVENT®

FOR ANIMAL TREATMENT ONLY KEEP OUT OF REACH OF CHILDREN

READ LABEL BEFORE USE

Directions for use:

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

For use in healthy broiler chickens one day of age or older.

Shake vial well before using. Dilute vaccine with clean tap water using a clean container. For spray cabinet application, include sterile green dye in the dilution to encourage oral ingestion.

SPRAY CABINET APPLICATION: Dilute each 10,000 dose vial according to the spray cabinet manufacturer's recommendations. When using a Huvepharma supplied spray cabinet, it is recommended to dilute 10,000 doses to 2,500mL.

ON FEED APPLICATION: Dilute each 10,000 dose vial to 4,000mL and apply to the first feed chicks receive. A method of application should be chosen that ensures rapid, even coverage of the total surface area of the feed available to the chicks, e.g. coarse spray over ready laid feed. Stir contents of sprayer periodically. Ensure a dose rate of 0.4mL per chick will be delivered based on expected feed consumption rate.

Precautions:

Feed or water containing anti-coccidial medicines should not be given to chicks receiving this vaccine for 21 days post-vaccination.

Withholding Period - Nil

58111010.02

ADVENT®

Registered and distributed by: Agri-Health NZ Ltd
0800 821 421
www.agrihealth.co.nz

AgriHealth
Evidence based vet medicines

Manufactured in the USA.
Huvepharma, Inc.
Lincoln, NE 68528
U.S. Veterinary License No. 605

HUVEPHARMA

Advent is registered pursuant to the ACVM Act 1997, No. A11395. See www.foodsafety.govt.nz for registration conditions.

FOR ANIMAL TREATMENT ONLY KEEP OUT OF REACH OF CHILDREN

ADVENT®

Coccidiosis vaccine containing live oocysts of *Eimeria maxima*, *E. tenella* and *E. acervulina*

10 X 10,000 Dose Vials

AgriHealth
Evidence based vet medicines

An aid in the prevention of coccidiosis in broiler chickens caused by *Eimeria maxima*, *E. tenella* and *E. acervulina*.

Store at 2-8°C away from sunlight. Do not freeze. Discard all unused concentrate and prepared product within 10 hours of vial opening.

Disposal: Preferably dispose of product by use. Dispose of empty containers, packaging and unused product in garbage.