

# Melovem 30

Meloxicam 30mg/mL

## READ CARTON LABEL AND LEAFLET BEFORE USE

**Directions for use:** Administer in combination with antibiotic therapy as appropriate.

**CATTLE: Administer 1mL per 60kg bodyweight** (0.5 mg meloxicam/kg) by subcutaneous or intravenous injection. For very young calves a dose rate of 0.3mL per 20kg is appropriate. For single use only.

**SHEEP: Administer 1mL per 30kg bodyweight** (1.0 mg meloxicam /kg) by subcutaneous injection high on the neck behind the ear. For single use only.

**PIGS: Administer 1mL per 75kg bodyweight** (0.4 mg meloxicam/kg) by intramuscular injection in the anterior neck. Can be repeated once after 24 hours if necessary.

**HORSES: Administer 1mL per 50kg bodyweight** (0.6 mg meloxicam/kg) by intravenous injection. For single use only.

Manufactured in Europe by **Dopharma**

Registered in New Zealand to:

**AgriHealth NZ Ltd. Freephone 0800 821 421**  
**www.agrihealth.co.nz**

Melovem 30 is registered pursuant to the  
ACVM Act 1997, No. A11562

See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

**AgriHealth**

**DANGER** May be harmful if swallowed. Causes skin irritation. Causes serious eye irritation. May cause an allergic skin reaction. Suspected of damaging fertility or the unborn child. May cause damage to organs through prolonged or repeated exposure. Harmful to terrestrial vertebrates.

**Handling precautions:** Do not handle until all safety precautions have been read and understood (refer leaflet inside carton).

**First Aid:** If medical advice is needed, have product container or label at hand. If exposed or concerned, or if you feel unwell, seek medical advice, or call the NATIONAL POISONS CENTRE 0800 POISON (0800 764 766).

**IF ON SKIN:** Wash with plenty of soap and water.  
**IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If skin rash or eye irritation occurs or persists, get medical advice/attention.

**Disposal:** Preferably dispose of product by use. Dispose of empty containers, packaging and unused product at an approved landfill or facility.

### Withholding Period:

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.**

**Meat:** Cattle -10 days Sheep - 11 days Pigs - 3 days  
**Horses - 28 days Milk:** Cattle and sheep - 84 hours  
Store below 25°C. Use within 90 days of opening.

# Melovem 30



Meloxicam 30mg/mL 100mL

**RESTRICTED VETERINARY MEDICINE  
KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY**

# Melovem 30



Meloxicam 30mg/mL 100mL

### Meloxicam 30mg/mL

Non-steroidal anti-inflammatory, analgesic and antipyretic injection for use in cattle, sheep, pigs and horses.

**Cattle:** For use in acute mastitis, in combination with antibiotic therapy, as appropriate, to reduce clinical symptoms in lactating cows. For use in acute respiratory infection with appropriate antibiotic therapy to reduce clinical signs. For use in diarrhoea in combination with oral re-hydration therapy to reduce clinical signs in calves over one week of age. To assist in the control of pain following the dehorning of cattle particularly that following heat cauterly dehorning of young cattle. It is recommended that the injection be administered approximately 10 minutes before dehorning and be accompanied by corneal nerve block anaesthesia.

**Sheep:** For use in sheep and lambs 14 days of age or older for the alleviation of pain and inflammation.

**Pigs:** For use in acute non-infectious locomotor disorders to reduce the symptoms of lameness and inflammation. For use in puerperal septicaemia and toxæmia (mastitis-metritis-agalactia syndrome) with appropriate antibiotic therapy.

**Horses:** For rapid initiation of therapy of musculoskeletal disorders and relief of pain associated with colic.

**AgriHealth**

58000343-4



EXP:

Lot: AgriHealth