

RESTRICTED VETERINARY MEDICINE
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

ALBAC® G Feed Supplement
Zinc Bacitracin

For the prevention and treatment of Necrotic Enteritis (*Clostridium perfringens* types A and C) in pigs and poultry.

ACTIVE DRUG INGREDIENT

Each kilogram contains feed grade zinc bacitracin equivalent to 150 grams of bacitracin (1 gram bacitracin is equivalent to 42,000 International Units).

Net contents: 25kg

MIXING DIRECTIONS

ALBAC G Feed Supplement should be premixed with 10-20 kg of feed and then thoroughly mixed with the bulk feed.

DOSAGE

	Kg ALBAC / tonne feed	Equivalent to bacitracin ppm	Indications for use
Broiler Chickens	0.33	50	As an aid in prevention of necrotic enteritis caused by or complicated by <i>Clostridium</i> spp. or other organisms susceptible to zinc bacitracin,
	1.33	200	As an aid in treatment of necrotic enteritis caused by or complicated by <i>Clostridium</i> spp. or other organisms susceptible to zinc bacitracin. CAUTION: To control a necrotic enteritis outbreak, start medication at the first clinical signs of disease. Administer continuously for 5-7 days or as long as clinical signs persist and then reduce medication to preventative level (0.33 kg ALBAC/tonne feed).
Quail	1.33	200	For the prevention of ulcerative enteritis in growing quail due to <i>Clostridium colinum</i> susceptible to zinc bacitracin. Feed continuously as the sole ration.
Pregnant Sows	1.65	250	For control of bacterial enteritis caused by <i>Clostridium perfringens</i> in suckling piglets when fed to sows from 14 days before through to 21 days after farrowing on premises with a history of clostridial scours. CAUTION: Diagnosis should be confirmed by a veterinarian when results are not satisfactory. Feed containing an approved level of zinc bacitracin should be the sole ration.

Indiscriminate use may increase the risk of the development of antibiotic resistance. Duration of use should be limited to risk periods when possible.

WITHHOLDING PERIOD: Nil

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WARNING: May cause an allergic skin reaction.

PRECAUTIONS: Read label before use. Avoid breathing dust. Contaminated work clothing should not be allowed out of the workplace. Wash contaminated clothing before reuse. Wear protective gloves, clothing and face protection.

STORAGE: Store below 30°C (Room Temperature) in a dry place. Close container securely after use.

DISPOSAL: Shake and empty containers into medicated feed. Do not dispose of undiluted chemicals on-site. Puncture or shred and bury empty container below 500 mm in a disposal pit specifically marked and set up for the purpose, clear of waterways, vegetation and roots. Empty bag and left-over product should not be burnt.

FIRST AID: IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs, seek medical advice/attention.

Registered pursuant to the ACVM Act 1997, No. A6736
See www.foodsafety.govt.nz for registration conditions

Approved pursuant to the HSNO Act 1996, No. HSR007846
See www.epa.govt.nz for approval conditions

Registered to: AgriHealth NZ Ltd www.agrihealth.co.nz

AgriHealth
Evidence based vet medicines

BATCH:

EXPIRY DATE: