

**RESTRICTED VETERINARY MEDICINE
FOR ANIMAL TREATMENT ONLY
CRITICALLY IMPORTANT ANTIBIOTIC**

Pharmasin[®] 25%

Granular Premix

TYLOSIN PHOSPHATE

FOR USE IN CHICKENS AND PIGS

Net contents: 20 kg

Active Ingredient: TYLOSIN (as TYLOSIN PHOSPHATE) 250 g/kg

Tylosin contains not less than 80% factor A

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT

INDICATIONS:

Broiler Chickens and Replacement Layers:

As an aid in the treatment and prevention of chronic respiratory disease caused by susceptible strains of *Mycoplasma gallisepticum* and/or *Mycoplasma synoviae*. For the treatment and prevention of necrotic enteritis associated with susceptible strains of *Clostridium perfringens*.

Pigs:

For the treatment and prevention of ileitis caused by *Lawsonia intracellularis*.

MIXING AND FEEDING DIRECTIONS:

IMPORTANT: MUST BE THOROUGHLY MIXED IN FEEDS BEFORE USE.

Animal Species	Recommended use	Feeding period	Weight of Pharmasin 25% Granular Premix per 1000 kg of Complete Feed	Use Level
Pigs	For the treatment and prevention of ileitis caused by <i>Lawsonia intracellularis</i>	Treatment - Feed continuously for 21 days Prevention - length of treatment required will depend on the level of disease risk and exposure, up to 21 days	400 g 160 g	100 ppm 40 ppm
Broiler Chickens and Replacement Layers	As an aid in the treatment and prevention of chronic respiratory disease caused by susceptible strains of <i>Mycoplasma gallisepticum</i> and/or <i>Mycoplasma synoviae</i> .	For 5 days after hatching. Repeat for 2 days at 3 to 5 weeks of age	3600 g to 4000 g	900 ppm to 1000 ppm
	For the treatment and prevention of necrotic enteritis associated with susceptible strains of <i>Clostridium perfringens</i> .	Treatment-treat for 7 days Prevention-feed during periods of risk of clinical disease	400 g 400 g	100 ppm 100 ppm
Layers	As an aid in the control of chronic respiratory disease caused by susceptible strains of <i>Mycoplasma gallisepticum</i>	Feed from time of shifting into laying house	88 g to 220 g	22 ppm to 55 ppm

Tylosin is an antibiotic in the macrolide family, and is considered critically important to human and animal health. Indiscriminate use of this antibiotic can contribute to the development of antibiotic resistance, and it should not be used as a first line treatment. The use of this antibiotic should be reserved for the treatment of conditions that respond poorly to other classes of antibiotic following culture and sensitivity testing, or in cases where other treatments have been clinically determined to be ineffective. Use of this antibiotic should be limited to the minimum period needed to meet the clinical objective. Clinical response should be monitored during treatment, and choice of therapy reviewed if clinical signs of disease persist, increase, or relapse.

WITHHOLDING PERIODS:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Meat: Pigs and poultry producing meat or offal for human consumption must not be sold for slaughter either during treatment or within **12 hours** of the last treatment.

Eggs: Nil

DANGER

Causes serious eye irritation. May cause allergy or asthma symptoms or breathing difficulties if inhaled.

Very toxic to aquatic life with long lasting effects.

Handling precautions: Wash hands and exposed skin thoroughly after handling. Contaminated work clothing should not be allowed out of the workplace. Wash contaminated clothing before reuse. Avoid breathing dust. In case of inadequate ventilation wear respiratory protection.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If eye irritation persists: get medical advice/attention.

IF INHALED: If breathing is difficult, remove to fresh air and keep at rest in a

position comfortable for breathing. If experiencing respiratory symptoms: Call the National Poisons Centre 0800 764 766 or a doctor or physician. **IF ON SKIN:** Wash with plenty of soap and water. If skin irritation or rash occurs: Get medical advice/attention.

Environmental Protection:

Avoid release to the environment of product or empty container. Collect spillage.

Disposal:

Preferably dispose of product by use. Dispose of product and packaging at an approved landfill or equivalent facility.

UN 3077 ENVIRONMENTALLY HAZARDOUS SUBSTANCE, SOLID, N.O.S. (tylosin phosphate 25%)

STORAGE: Store in original pack, in a dry place, below 25°C. Protect from light. Do not refrigerate or freeze - protect from frost.

Restricted Veterinary Medicine. Registered pursuant to the ACVM Act 1997, No. A010451
See www.foodsafety.govt.nz for registration conditions.

Batch:

Expiry:



3 800043 755359

Manufactured by:


Huvepharma E00D

www.huvepharma.com

Registered to: AgriHealth NZ Ltd

www.agrihealth.co.nz,

Phone 0800 821 421



AgriHealth



LB677v6-Phar25-mp20kg-NZ0726