

BioBos RCC

Vaccine for cattle containing inactivated Bovine Rotavirus, Bovine Coronavirus, and *E.coli* expressing F5 (K99) adhesin. Contains thiomersal (0.01% w/v).



492/K/125d/NZ/1

RESTRICTED VETERINARY MEDICINE
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

BioBos RCC



Vaccine for cattle containing inactivated Bovine Rotavirus, Bovine Coronavirus, and *E.coli* expressing F5 (K99) adhesin.

Contains thiomersal (0.01% w/v).

Net Contents 250mL (125 doses)

Handling precautions:

Wear protective gloves when handling. Wash exposed skin thoroughly after handling.

First Aid:

If exposed or concerned or if you feel unwell: Get medical advice or attention. For advice call the National Poisons Centre 0800 POISON (0800 764 766).

Disposal:

Any unused product or empty containers should be disposed of at an approved landfill or other approved facility.

Store between 2 - 8°C. Do not freeze. Protect from light. Use within 10 hours of opening.

BioBos RCC is registered pursuant to the ACVM Act 1997, No. A12072

See www.foodsafety.govt.nz for registration conditions.


Manufactured in Europe by Bioveta a.s., Ivanovice na Hané, Czech Republic. www.bioveta.cz

Batch:

Expiry:

RESTRICTED VETERINARY MEDICINE
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

BioBos RCC



Vaccine for cattle containing inactivated Bovine Rotavirus, Bovine Coronavirus, and *E.coli* expressing F5 (K99) adhesin.

Contains thiomersal (0.01% w/v).

Net Contents 250mL (125 doses)

READ LABEL AND LEAFLET BEFORE USE

Indications

For active immunisation of pregnant cows and heifers to stimulate the development of antibodies against Bovine Rotavirus, Bovine Coronavirus, and *E.coli* expressing F5 (K99) adhesin, and to increase the level of passive immunity of calves against neonatal diarrhoea caused by rotavirus, coronavirus and enteropathogenic *E.coli*. When calves consume colostrum and milk from vaccinated cows during the first two weeks of life, these antibodies prevent neonatal diarrhoea caused by bovine rotavirus, coronavirus and *E.coli* expressing F5 (K99) adhesin, and reduce faecal shedding of virus in calves infected with bovine rotavirus and coronavirus.

Directions for use

Shake well before use.

CATTLE: Administer 2 mL, by intramuscular injection 12-3 weeks prior to calving. Annually revaccinate with a single dose of BioBos RCC 12-3 weeks prior to each subsequent calving.

Precautions:

Do not vaccinate unhealthy animals.

Withholding Period: Nil

AgriHealth

Registered to: AgriHealth NZ Ltd
Phone 0800 821 421 www.agrihealth.co.nz

