

Vaccine for cattle containing
inactivated Bovine herpesvirus
type 1 (BoHV-1) Contains thiomersal

Batch:
Expiry:


Biobos IBR
MARKER VACCINE



184/K/100ml/NZ/3

**RESTRICTED VETERINARY MEDICINE
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY**

Biobos IBR
MARKER VACCINE



Vaccine for cattle containing
inactivated Bovine herpesvirus
type 1 (BoHV-1). Contains thiomersal

Net Contents 100mL (50 doses)

WARNING

May cause an allergic skin reaction.
Suspected of causing cancer.

Handling precautions:

Obtain special instructions before use.
Do not handle until all safety precautions
have been read and understood.
Wear gloves when handling. Wash
exposed skin thoroughly after handling.
Avoid breathing mist or spray.
Contaminated work clothing should not
be allowed out of the workplace. Wash
contaminated clothing before reuse.
Store locked up.

First Aid:

IF ON SKIN: Wash with plenty of soap and
water. If skin irritation or rash occurs, or if
exposed or concerned: get medical
advice or attention.

For advice call the National Poisons
Centre 0800 POISON (0800 764 766)

Disposal:

Any unused product or empty containers
should be disposed of at an approved
landfill or other approved facility.

Registered pursuant to the ACVM
Act 1997, No. A11239.


See www.foodsafety.govt.nz
for registration conditions.

Manufactured in Europe by Bioveta, a.s.,
683 23 Ivanovice na Hané, Czech Republic
www.bioveta.cz



**RESTRICTED VETERINARY MEDICINE
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY**

Biobos IBR
MARKER VACCINE



Vaccine for cattle containing
inactivated Bovine herpesvirus
type 1 (BoHV-1). Contains thiomersal

Net Contents 100mL (50 doses)

READ LABEL AND LEAFLET BEFORE USE

Indications

For active immunisation of cattle to
reduce intensity and duration of the
clinical symptoms caused by infection
with the Bovine herpesvirus type-1 (IBR)
virus and to reduce excretion of the
field virus.

Directions for use

Shake well before use.

CATTLE: Administer 2 mL, by
intramuscular injection into the anterior
half of the neck.
Initial vaccination should be given at
three months of age or older to avoid
interference by maternal antibodies.
Give booster vaccination three weeks
after first (sensitiser) vaccination.

Withholding Period - Nil

Store between 2 - 8°C. Do not freeze.
Use within 28 days of opening.
Protect from light.

Registered in New Zealand to:

AgriHealth NZ Ltd
0800 821 421
www.agrihealth.co.nz

AgriHealth



8 594004 859761