



**KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY**

# Coxipol<sup>®</sup> 25%

## microGranulate Premix for Chickens

**Net contents: 20 kg**

**Active ingredient:** CLOPIDOL 250 g/kg

**READ THE ENTIRE LABEL BEFORE USE**

**INDICATIONS:**

As an aid in the prevention of coccidiosis caused by *Eimeria acervulina*, *E. maxima*, *E. brunetti*, *E. necatrix* and *E. tenella* in broiler chickens.

**DIRECTIONS FOR USE:**

**By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.**

**POULTRY:**

For use in broiler chickens. Coxipol<sup>®</sup> 25% should be mixed into a secondary premix prior to incorporation in finished feed. Add 0.5kg of Coxipol<sup>®</sup> 25% to each tonne of chicken feed. This produces a clopidol content of 125 g/tonne in the finished feed which should be fed continuously as the sole ration.

**WITHHOLDING PERIOD:**

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.**

**Meat:** Animals producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 2 days of the last treatment.

**Eggs:** Not for use in animals from which eggs are produced for human consumption.

**STORAGE:** Store below 30°C in a dry place.

**WARNING**

May cause damage to organs through prolonged or repeated exposure.  
This substance must only be used as a veterinary medicine administered in poultry feed.

**Handling precautions:**

Wear protective gloves, face protection and coveralls buttoned to the neck when handling.  
Wash hands and exposed skin thoroughly after handling.  
Take off contaminated clothing and wash before reuse. Do not breathe dust.

**First Aid:**

Get medical advice if you feel unwell.  
For advice see a doctor, or call the National Poisons Centre 0800 POISON (0800 764 766)

**Disposal:**

Preferably dispose of product by use. Shake empty bag into medicated feed. Dispose of unused product and empty packaging at an approved landfill or other facility.



**Manufactured in Europe by:** HUVEPHARMA EOOD  
[www.huvepharma.com](http://www.huvepharma.com)

®Trademark registered by HUVEPHARMA EOOD

Registered pursuant to the ACVM Act 1997, No A11356.  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

**Registered to:** AgriHealth NZ Ltd, [www.agrihealth.co.nz](http://www.agrihealth.co.nz), Phone 0800 821 421

**AgriHealth**

**BATCH:**

**EXPIRY:**

