

70/148 mm

Pantone 313 C

Pantone 377 C

Pantone 355 C

Batch  
Expiry



LB1016v5Phar200inj250-NZ0720



FOR ANIMAL TREATMENT ONLY  
KEEP OUT OF REACH OF CHILDREN  
RESTRICTED VETERINARY MEDICINE

## TYLOVET INJECTION<sup>®</sup>

Tylosin 200mg/mL  
solution for injection

250 mL

For intramuscular use in cattle,  
pigs, sheep and goats.  
For slow intravenous use in cattle.



### TYLOVET INJECTION<sup>®</sup>

Read enclosed insert prior to use.

**DOSAGE:**

**Cattle:** 5 to 10 mg/kg (2.5 - 5 mL / 100kg)  
liveweight daily  
**Pigs:** 5 to 10 mg/kg (0.25 - 0.5 mL / 10kg)  
liveweight daily  
**Sheep and Goats:** 10 mg/kg (0.5 mL / 10kg)  
liveweight daily

**INDICATIONS:**

**Cattle:** Treatment and control of acute mastitis, metritis, respiratory infections, foot-rot (necrotic pododermatitis), calf diphtheria.  
**Pigs:** Treatment and control of swine dysentery and enteritis associated with the presence of *Campylobacter coli* and other organisms sensitive to tylosin, swine erysipelas, pneumonia and arthritis due to *Mycoplasma*.  
**Sheep and Goats:** Treatment of the early stages of peracute and acute contagious agalactia caused by *Mycoplasma agalactiae* and caprine pleuropneumonia caused by *Mycoplasma mycoides var capri (M capri)*.  
The prophylactic and therapeutic use of this product should only be for the minimum period needed to meet the clinical objective. Indiscriminate use of this product could contribute to antibiotic resistance.

Produced in Europe by Huvepharma NV  
Registered to and distributed by AgriHealth NZ Ltd  
Restricted Veterinary Medicine  
Registered pursuant to the ACVM Act 1997,  
No A010807. See [foodsafety.govt.nz](http://foodsafety.govt.nz) for registration conditions  
Approved pursuant to the HSN0 Act 1996, NO  
HSR002368



Solution for injection  
Cattle, sheep, goats, swine



FOR ANIMAL TREATMENT ONLY  
KEEP OUT OF REACH OF CHILDREN  
RESTRICTED VETERINARY MEDICINE

## TYLOVET INJECTION<sup>®</sup>

Tylosin 200mg/mL  
solution for injection

250 mL

For intramuscular use in cattle,  
pigs, sheep and goats.  
For slow intravenous use in cattle.



### TYLOVET INJECTION<sup>®</sup>

**WARNING**

May cause an allergic skin reaction

Causes eye irritation

Safety precautions and directions

Wear protective gloves. Wash hands thoroughly after handling. Contaminated clothing should not be allowed out of the workplace. Wash contaminated clothing before reuse.

First Aid

**If in the eyes:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists, get medical advice.

**If on skin:** Wash with plenty of soap and water. If skin irritation or rash occurs, get medical advice. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor.

**WITHHOLDING PERIODS:**

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

**Meat:** Animals producing meat and offal for human consumption must not be sold for slaughter either during treatment or within **21 days** of the last treatment.

**Milk:** Milk intended for sale for human consumption or manufacture for human consumption must be discarded during treatment and for **72 hours** following the last treatment.

STORE below 25° C, protect from light.  
Discard unused content 28 days after opening.



Solution for injection  
Cattle, sheep, goats, swine