


FOR ANIMAL TREATMENT ONLY KEEP OUT OF REACH OF CHILDREN

ToltraVet Poultry 2.5%

FOR THE TREATMENT OF COCCIDIOSIS IN POULTRY



Net contents: 1 litre
25g/L toltrazuril

AH-TVP.26

AgriHealth
Evidence based vet medicines

Dosage and administration: By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

For oral administration in drinking water. Ensure all birds have access to medicated water. ToltraVet Poultry 2.5% should be mixed with water in the water reservoir, and is not recommended for use by predilution or administration via proportional dosing pumps.

Dose rate: 7mg of toltrazuril (0.28 mL ToltraVet Poultry 2.5%) per kg body weight per day for 2 consecutive days. Multiply this dose per bird by the total number of birds to be treated.

For soft water: add the calculated dose to the total drinking water volume consumed per day for continuous treatment over a 24 hour period. Refresh medicated water after the first 24 hours.

For hard water: add the calculated dose to the drinking water consumed over an 8 hour period, then replace with unmedicated water for the following 16 hour period. Repeat on the second day.

	Body weight per bird			Volume ToltraVet per day		
	10 birds	100 birds	1000 birds			
1.0 kg	2.8 mL	28 mL	280 mL			
1.5 kg	4.2 mL	42 mL	420 mL			
2.0 kg	5.6 mL	56 mL	560 mL			

The dose rate should be checked against water intake and varied accordingly. When adding ToltraVet Poultry 2.5% to water, avoid foam formation. Although foam formation does not affect the made up concentration, it may temporarily impair correct functioning of various parts in the drinking water systems.

Withholding period

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Meat: Chickens producing meat and offal for human consumption must not be sold for slaughter during or within 14 days of the last treatment.

Eggs: Eggs from treated birds should not be sold for human consumption 10 days following cessation of last treatment.

WARNING

READ LABEL BEFORE USE
Suspected of damaging fertility, or the unborn child.
May cause damage to organs through prolonged or repeated exposure.
Toxic to aquatic life with long lasting effects.



Handling precautions:

Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Wear gloves, eye and face protection when handling. Wash exposed skin thoroughly after handling.

Do not breathe mist or spray. Store locked up

First Aid:

If exposed or concerned, or if you feel unwell: Get medical advice or attention.

For advice call the National Poisons Centre
0800 POISON (0800 764 766).

Environmental protection:

Avoid release to the environment. Collect spillage.

Disposal: Any unused product or empty containers should be disposed of at an approved landfill or other approved facility.

Storage:

Store below 25°C.

ToltraVet Poultry 2.5% is registered pursuant to the ACVM Act 1997, No. A11267.

See www.foodsafety.govt.nz for registration conditions.

Manufactured by ALKE Healthcare Products, Turkey.

Registered in New Zealand to:

Agrihealth NZ Ltd
Phone 0800 821 421
www.agrihealth.co.nz

AgriHealth

T-02



BATCH:

EXPIRY: