

Warnings

Do not use concurrently with another non-steroidal anti-inflammatory. Do not administer where there is abundant haemorrhage. Do not use in animals suffering from cardiac, hepatic or renal disease. Do not administer by the intra-arterial route. Do not administer to an animal that has developed hypersensitivity to ketoprofen. Abomasal ulceration and perforation has been seen when used in calves at disbudding.

Handling Precautions

KEEP OUT OF REACH OF CHILDREN

WARNING: Harmful if swallowed. Causes skin and eye irritation. Harmful to terrestrial vertebrates. Wash hands and exposed skin thoroughly after handling. Do not eat, drink or smoke when using this product. Wear protective gloves. Take off contaminated clothing and wash before reuse. If medical advice is needed, have product container or label at hand.

IF SWALLOWED: Call the National Poisons Centre 0800 764 766 or a doctor/physician if you feel unwell. Rinse mouth.

IF ON SKIN: Wash with plenty of soap and water. If skin irritation occurs: Get medical advice / attention.


IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.

DISPOSE of empty container by wrapping with paper and putting in the garbage. Avoid release of the product into the environment.

KetoMax 15% is registered pursuant to the ACVM Act 1997, No. A11031. See www.foodsafety.govt.nz for registration conditions.

RESTRICTED VETERINARY MEDICINE FOR ANIMAL TREATMENT ONLY

KetoMax 15%



For use in cattle for the anti-inflammatory, analgesic and anti-pyretic treatment of:

- Acute mastitis
- Mammary oedema - Colic
- Respiratory conditions - Musculoskeletal conditions

READ THE LABEL BEFORE USING THIS PRODUCT

Directions for Use

CATTLE: Administer by intravenous or intramuscular injection, in the neck. DOSE: 1 mL / 50 kg / day (3 mg/kg ketoprofen) for 1 to 3 days depending on severity and duration of symptoms

Properties

Ketoprofen is a non-steroidal anti-inflammatory drug (NSAID) of the propionic acid class. The anti-inflammatory, analgesic and anti-pyretic effects of ketoprofen are attributed to the inhibition of cyclooxygenase in the inflammatory process. Ketoprofen is rapidly and extensively absorbed via either intramuscular or intravenous injection.

Withholding Periods

Meat: Cattle producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 2 days of the last

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Store below 25°C (room temperature). Use remainder of vial within 90 days of opening, or discard unused portion.

Registered to AgriHealth NZ Ltd Freephone 0800 821 421 www.agrihealth.co.nz

ecuphar®

ecuphar

AgriHealth

40035597

Batch

Expiry



GTIN: 08431506341717

CTN KetoMax 15% 250ml NZ Date 29/07/2021 EX2888-21-01 C661 69x69x149 mm / Control Lines 1x4: 45mm 2x5: 6