



**DANGER**  
**KEEP OUT OF REACH OF CHILDREN**  
**FOR ANIMAL TREATMENT ONLY**



# Sacox<sup>®</sup> 120

## microGranulate Premix for Poultry & Pigs

### NET CONTENTS: 25kg

BY LAW, THIS PRODUCT MUST NOT BE USED FOR THE PURPOSE OF  
 PROMOTING GROWTH OR INCREASED YIELD IN RUMINANTS.

**Active Ingredients:** SODIUM SALINOMYCIN 120 g/kg  
**Net contents:** 25 kg net

**READ THE ENTIRE LABEL BEFORE USE**

#### INDICATIONS:

For the prevention of coccidiosis caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. mivati*, *E. necatrix* and *E. tenella* in broiler chickens and in replacement birds intended for use as caged layers. For enhancing productivity by increasing the rate of weight gain and improving feed efficiency of grower/finisher pigs.

#### CONTRAINDICATIONS & CONSTRAINTS:

DO NOT exceed recommended use levels.  
 DO NOT feed to horses or any other animals or birds not included in the label claim. Salinomycin can cause cardiac and muscular paralysis and conduction defects. Some species, such as horses and dogs, are particularly susceptible. Deaths from salinomycin may occur in horses and dogs.  
 DO NOT feed to animals receiving tiamulin. Tiamulin inhibits biotransformation in the liver, thus stopping detoxification. Other drugs, especially antibiotics such as erythromycin and tylan (macrolides) and lincomycin and clindamycin (lincosamides) have this property. Delaying elimination of salinomycin may necessitate a meat withholding period. Care must be exercised when feeding concurrently with other anti-microbials.

#### DIRECTIONS FOR USE

By law, the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as recommended on the label.

#### POULTRY

For use in broiler chickens and replacement pullets only. Sacox<sup>®</sup> 120 should be mixed into a secondary premix prior to incorporation in finished feed. Add 0.5kg of Sacox<sup>®</sup> 120 premix to each tonne of chicken feed. This produces a salinomycin level of 60 g/tonne in the finished feed which should be fed continuously as the sole ration.

NOTE: Medicated feed must be labelled with appropriate precautions.

#### PIGS

For piglets less than 4 months of age: add 0.25 to 0.5 kg of Sacox<sup>®</sup> 120 to each tonne of feed. This produces a salinomycin level of 30 to 60 g/tonne in the finished feed.

For pigs greater than 6 months of age: add 0.125 to 0.25 kg of Sacox<sup>®</sup> 120 to each tonne of feed. This produces a salinomycin level of 15 to 30 g/tonne in the finished feed. Sacox<sup>®</sup> 120 should be mixed into a secondary premix prior to incorporation in finished feed.

NOTE: Medicated feed must be labelled with appropriate precautions.

Registered pursuant to the ACVM Act 1997, No. A007640  
 See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

Registered to: AgriHealth NZ Ltd

[www.agrihealth.co.nz](http://www.agrihealth.co.nz)

Phone 0800 821 421

Manufactured By:

HUVEPHARMA FOOD

[www.huvepharma.com](http://www.huvepharma.com)

**AgriHealth**



3 800043 705675

#### DANGER

Toxic if swallowed  
 Causes skin irritation  
 May cause damage to the nervous system through prolonged or repeated exposure  
 Causes serious eye damage  
 Hazardous to terrestrial vertebrates



#### Handling Precautions

Avoid contact with eyes and skin.  
 Wear protective gloves, clothing, eye and face protection. Do not breathe dust.  
 Do not eat, drink or smoke while using this product.  
 Remove protective clothing and wash before reuse.  
 Wash hands and exposed skin thoroughly after handling.  
 Store product locked up.

#### First Aid

If medical advice is needed, have product container or label at hand.

IF SWALLOWED: Immediately call the National Poisons Centre (0800 764 766) or doctor/physician. Do NOT induce vomiting. Rinse mouth.

IF ON SKIN: Wash with plenty of soap and water

If skin irritation occurs: get medical advice/attention

If exposed or if you feel unwell: Call the National Poisons Centre or a doctor/physician.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. Immediately call the National Poisons Centre or a doctor/physician.

Symptoms of poisoning may include irregular breathing, tremor, and eye and skin irritation.

#### Environmental Protection:

Hazardous to terrestrial vertebrates. Avoid release to the environment. Collect spillage.

In case of spillage wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material.

**Disposal:** Shake empty bag into medicated feed. Destroy empty containers by puncturing or shredding. Dispose of product, waste and packaging at an approved landfill or equivalent facility.

#### WITHHOLDING PERIODS:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

**EGGS:** Eggs from treated birds must not be sold for human consumption for 10 days following the last treatment.

**MEAT:** Nil.

**STORAGE:** Store below 25°C (room temperature) in a dry place.

Batch:

Expiry:



C/PAP/PE